

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
21 April 2005 (21.04.2005)

PCT

(10) International Publication Number
WO 2005/035742 A2

- (51) International Patent Classification⁷: C12N 5/10, A61K 35/00
- (21) International Application Number: PCT/US2004/033220
- (22) International Filing Date: 7 October 2004 (07.10.2004)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
- | | | |
|------------|-----------------------------|----|
| 60/509,928 | 9 October 2003 (09.10.2003) | US |
| 60/510,021 | 8 October 2003 (08.10.2003) | US |
| 60/510,022 | 8 October 2003 (08.10.2003) | US |
| 60/510,072 | 9 October 2003 (09.10.2003) | US |
- (71) Applicant (for all designated States except US): VET-STEM INC. [US/US]; 12120 Tech Center Drive, Suite D, Poway, California 92064 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): HARMAN, Robert, J. [US/US]; 1460 Cedar Street, Ramona, California 92065 (US). SAND, Theodore, T. [US/US]; 12745 Corte Reyito, Poway, California 92064 (US).
- (74) Agents: LAHERTY, Carol, D.? et al.; Seed Intellectual Property Law Group PLLC, 701 - 5th Avenue, Suite 6300, Seattle, Washington 98104-7092 (US).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: METHODS OF PREPARING AND USING NOVEL STEM CELL COMPOSITIONS AND KITS COMPRISING THE SAME

(57) Abstract: The present invention provides novel stem cell compositions having significant therapeutic and practical advantages, as well as methods of preparing and using such compositions for the treatment and prevention of injury and disease in patients. The invention may be applied to stem cell populations isolated from a wide variety of animals, including humans, and tissues. In particular applications, the invention is used to prepare a stem cell composition from a collagen-based tissue, such as adipose tissue, isolated from a patient, and the stem cell composition is subsequently provided to a site of actual or potential injury in the patient. The invention further includes related kits comprising the stem cell compositions, which are remarkably stable and retain viability and efficacy during storage and shipment.



WO 2005/035742 A2

METHODS OF PREPARING AND USING NOVEL STEM CELL COMPOSITIONS AND KITS COMPRISING THE SAME

BACKGROUND OF THE INVENTION

5 Field of the Invention

The present invention is generally related to the field of stem cell-based therapy and, more specifically, to compositions comprising stem cells and methods of preparing and using compositions comprising stem cells for the treatment and prevention of injury and disease.

10 Description of the Related Art

Due to their remarkable ability to regenerate and develop into a variety of cell types, stem cells possess great therapeutic potential in the treatment of a wide variety of diseases and injuries, particular those involving the destruction or damage of normal tissue, such as spinal cord injuries,
15 Parkinson's Disease, Alzheimer's Disease, and multiple sclerosis. Until fairly recently, it was thought that multipotent stem cells could only be isolated from embryonic tissue. However, it has now been discovered that multipotent stem cells exist in a variety of adult tissues, including bone marrow, skin, brain, muscle and adipose tissue. This discovery has sparked increased interest in
20 stem cell-based therapies, since such adult stem cells are more readily available than embryonic stem cells, and their use does not raise the same ethical concerns.

One significant limitation to the therapeutic use of stem cells is that they are present in very low numbers in most adult tissues, and their
25 isolation and purification is a tedious and expensive process. The generally practiced method of preparing stem cells before providing them to a patient involves purifying cells from a tissue sample, isolating stem cells, e.g., using

antibodies specific to stem cell surface markers from other cells, and/or culturing the cells. In addition, many procedures further involve treating the cells with an agent that induces differentiation down a specific lineage. The reagents used for the purification, culturing and differentiation of stem cells are very expensive, thus limiting the availability of stem cell-based therapies. In addition, the procedures involved in preparing and storing stem cells can result in cell death and loss of function, thus reducing the number of useful stem cells isolated and limiting their ability to be stored and shipped prior to use.

Clearly, there is a need in the art for improved methods of preparing stem cell populations suitable for therapeutic and prophylactic use, including methods capable of purifying a high number of viable cells from a tissue sample, particularly for autologous uses, which may be performed with increased ease and reduced costs. In addition, there is a related need for methods of preparing purified stem cells from a patient's tissue sample obtained by a medical professional, and providing the purified cells to a medical professional for administration to the patient. The present invention meets these needs by providing novel methods of preparing stem cell populations, novel compositions comprising stem cells, and streamlined procedures for preparing and providing stem cells for delivery to a patient.

BRIEF SUMMARY OF THE INVENTION

The present invention provides improved methods of preparing stem cells, improved compositions comprising stem cell populations, improved methods of treating and preventing injury or disease using the stem cell compositions, and kits comprising stem cell compositions.

In one embodiment, the invention includes a method of preparing a purified cell population comprising stem cells for introducing into a patient, comprising obtaining adipose tissue from said patient, processing said adipose tissue to separate cells therein from other tissue components, and purifying the separated cells from other tissue components.

In a related embodiment, the invention provides a method of preparing a composition comprising purified adipose tissue-derived stem cells for introducing into an animal, comprising obtaining adipose tissue from the tail head region of said animal, processing said adipose tissue to separate cells
5 therein from other tissue components, and purifying the separated cells from other tissue components.

In another embodiment, the invention includes a method of providing a composition comprising purified stem cells for introducing into a patient, comprising processing collagen-based tissue obtained from a patient to
10 separate cells therein from other tissue components, purifying the separated cells, and placing said separated cells in a container, thereby providing a composition comprising purified stem cells. In particular embodiment, the container is a syringe, vial, or cryovial.

In a further embodiment, the invention includes a method of
15 preparing a purified cell population comprising collagen-based tissue-derived stem cells for delivery to a patient, comprising obtaining collagen-based tissue from said patient, processing said collagen-based tissue to separate cells therein from other tissue components, and purifying the separated cells, wherein said processing comprises contacting said collagen-based tissue with
20 a series of screens, thereby preparing a purified cell population comprising collagen-based tissue-derived stem cells. In a further embodiment, the processing also includes treating the tissue with an enzyme that facilitates the release of cells from other tissue components.

In yet another related embodiment, the invention provides a
25 method of preparing a purified cell population comprising collagen-based tissue-derived stem cells for providing to a patient, comprising obtaining collagen-based tissue from said patient and processing said collagen-based tissue to isolate cells therein from other tissue components, wherein said processing comprises contacting the tissue with a surface to which the tissue
30 adheres, thereby preparing a purified cell population comprising collagen-based

tissue-derived stem cells. In a further embodiment, the processing also comprises mincing said tissue and/or treating the tissue with an enzyme that facilitates the release of cells from other tissue components. In one embodiment, contacting comprises mixing the tissue with particles of the
5 adherent surface.

In particular embodiments encompassing a surface to which the tissue adheres, the surface is Velcro, polystyrene, glass fiber, glass wool, cellulose, or ceramic.

In a related embodiment, the invention includes a device adapted
10 for preparing a cell population comprising collagen-based tissue-derived stem cells, comprising a series of mesh screens, arrayed such that one or more of the screens may be separated from each other while in contact with collagen-based tissue sample. In one embodiment, the screens comprise edges capable of cutting an adipose tissue sample. In another embodiment, the
15 device comprises a cutting implement that may be inserted between adjacent screens. In a further embodiment, the device further includes a container comprising said mesh screens, which may include an opening through which an adipose tissue sample may be placed into said container.

In a variety of embodiments, methods of the invention further
20 include suspending prepared cells in a physiologically compatible solution, and/or freezing or lyophilizing the cells.

In other embodiments, the methods further comprise shipping cells and compositions to a physician or veterinarian.

In particular embodiments of the invention, the collagen-based
25 tissue is adipose tissue or umbilical cord matrix.

In certain embodiments, methods of preparing a cell population or composition comprising stem cells further comprise suspending said purified cells in a physiologically compatible buffer, placing said purified cells in a syringe, and/or freezing cells in freezing medium.

In particular embodiments, methods of preparing a cell population or composition comprising stem cells do not include isolating stem cells from other purified cells.

In related embodiments, methods of processing tissue include
5 mincing the adipose tissue, treating the adipose tissue with an enzyme that facilitates the release of cells from other tissue components, exposing the adipose tissue to ultrasonic energy; and/or treating the adipose tissue with perfluorocarbons. In one embodiment, enzyme treatment is performed at a temperature below 28°C and/or a pH below 7.0.

10 The invention further includes a method of treating an injury or disease in a patient, comprising providing to said patient an isolated cell population or composition comprising collagen-based tissue-derived stem cells prepared according to a method of the invention.

Similarly, the invention includes a method of preventing an injury
15 in a patient, comprising providing to said patient an isolated cell population or composition comprising collagen-based tissue-derived stem cells prepared according to a method of the invention.

In particular embodiments, the injury or disease is a musculo-skeletal injury or disease. In certain embodiment, the injury is a sprain, strain,
20 dislocation, bruising, tear, or fracture. In other embodiments, the injury or disease is an ischemic injury or disease or a septic injury or disease. In various embodiments, the tissue is tendon, ligament, cartilage, or bone; hoof laminae; or lung, blood vessels, liver, nerve, or heart.

In certain embodiments, the isolated cell population or
25 composition is provided directly to a site of injury or disease. In another embodiment, the isolated cell population or composition is provided to the bloodstream of said patient. In one embodiment, the isolated cell population or composition is provided by injection, intravenously or inter-arterially.

In related embodiments, the methods of the invention are
30 practiced on a variety of different animals. Accordingly, in various

embodiments, cell populations are prepared from tissue isolated from any animal (including patients and donors), cell populations include cells prepared from tissue from any animal, and methods of treatment are performed on any animal. Accordingly, in particular embodiments, a patient or animal is a human, 5 a non-human animal, a horse or camel, a dog or cat, an exotic or zoological animal, a hoofed mammal, a bird, or a cow or goat. In various embodiments, the invention is practiced on patients or animals having significant commercial value, such as performance or racing animals, zoo animals, livestock or farm animals, dairy animals, companion animals, and rare or exotic animals.

10 In certain embodiments, tissue is obtained from any of a variety of sites on a patient or donor, including the tail head region. In other embodiments, tissue is a collagen-based tissue.

In various embodiments, compositions and cell populations, which comprise stem cells, further comprise one or more additional cellular or non- 15 cellular tissue component. In one embodiment, additional cells are blood cells, white blood cells, fibroblasts, fibroblast-like cells, neutrophils, monocyte/macrophages, and/or basophils. In another embodiment, additional tissue components are extracellular matrix polypeptides or fragments thereof, proteoglycans, cytokines, or growth factors. In one embodiment, an 20 extracellular matrix polypeptide is collagen, thrombospondin, fibronectin, vitronectin, cytotactin, laminin or an integrin.

In additional embodiments, the invention further includes kits comprising a cell population or composition of the invention. In one embodiment, the invention provides a kit useful in the treatment of an injury or 25 disease in an animal, comprising: a container comprising a composition comprising a stem cell population purified from a collagen-based tissue obtained from an animal.

In a related embodiment, the invention includes a kit useful in the prevention of an injury in an animal, comprising: a container comprising a

composition comprising a stem cell population purified from a collagen-based tissue obtained from an animal.

In yet another related embodiment, the kit comprises two or more containers, each comprising a composition comprising a stem cell population
5 purified from a collagen-based tissue obtained from an animal.

In one embodiment, a kit is useful in the treatment or prevention of a musculo-skeletal tissue injury.

In another kit embodiment, the stem cell population is present in a physiologically compatible solution. In various related embodiments, the
10 container is a syringe, a vial, or a cryovial. In further embodiments, the composition is frozen or lyophilized.

In certain kit embodiments, the collagen-based tissue was obtained from the animal to be treated. In one embodiment, the collagen-based tissue is adipose tissue obtained from the tail head region of the animal. In
15 other embodiment, the tissue is tendon, ligament, cartilage, or bone. In yet another embodiment, the tissue is lung tissue, blood vessels, liver, nerve, and heart. In another embodiment, the tissue is hoof laminae.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides novel and improved methods of
20 preparing stem cells useful for the therapeutic and prophylactic treatment of injuries and diseases. The invention further includes cell populations and compositions comprising stem cells purified according to such methods. The invention is based, in part, upon the surprising discovery that stem cell populations purified according to the methods of the invention have increased
25 efficacy and increased stability as compared to stem cells purified using previous methods, which typically include one or more additional steps of isolating stem cells from other cells present in a tissue source and/or culturing or differentiating the isolated stem cells prior to administration to a patient. Accordingly, the invention provides methods of preparing a stem cell population

that are simpler and more convenient to perform than prior methods.

Furthermore, the invention includes methods related to the preparation of convenient-to-use stem cell compositions that can be supplied in an applicator, such as a syringe, for administration to a patient.

5 A. Methods of Purifying Stem Cells

In general, the basic method of the invention includes processing a tissue sample comprising stem cells to separate cells therein from other tissue components and purifying the separated cells from other tissue components. In certain embodiments, the method does not include one or
10 more of the steps of isolating stem cells from other purified cells, culturing the purified cells, or differentiating the purified cells. In other embodiments, a method of the invention further comprises one or more of these steps.

The invention may be used to isolate stem cells from any tissue source, including but not limited to adipose tissue, umbilical cord matrix, brain
15 tissue, blood, muscle, bone marrow, tooth tissue and skin. In one embodiment, the tissue is a collagen-based tissue, such as adipose tissue or umbilical cord matrix. The methods of the invention are particularly well-suited to processing collagen-based tissues to facilitate the release of stem cells.

Tissue may be obtained from humans and other animals. In one
20 embodiment, the animal is a mammal. In certain embodiments, tissue is obtained from animals having significant commercial or economic value, such as horses, dogs, cats, camels, and cows. Similarly, in other embodiments, the tissue is obtained from performance or sport animals, such as horses and dogs. In other embodiments, tissue is obtained from a companion animal, such as a
25 dog or cat. In a further embodiment, tissue is obtained from exotic dogs (wolves, jackals, dingos, *etc.*) and exotic cats (leopard, fisher cat, mountain lion, jagarundi, tiger, lion, cheetah, *etc.*). In other embodiments, tissue is obtained from exotic or zoological animal, including but not limited to hoof stock, including ungulates or hoofed mammals, including, *e.g.*, gazelle, buffalo,

rhinoceros, eland, etc. In another embodiment, a tissue sample is obtained from a high value exotic species, including but not limited to birds, including, e.g., parrots, macaws, etc. In certain embodiments, tissue is obtained from large exotic or zoological animals, including but not limited to, zebras, lions, tigers, elephants, rhinoceroses, hippopotamuses, bears, and giraffes.

Tissue may be obtained from fetuses, or juvenile or adult animals, including humans. Furthermore, tissue may be obtained from the patient to whom the purified cells will be provided, or, alternatively, tissue may be obtained from a donor and the purified cells provided to a different patient. As such, the invention contemplates both autologous and allogeneic uses of the purified cells.

Tissue may be isolated from a patient or donor by any means available in the art. In certain embodiments, tissue is isolated by lipoaspiration, surgical removal, withdrawal using a needle and syringe, or lipectomy. A variety of additional procedures are described in U.S. Patent Application Publication No. 2003/0161816 A1 and U.S. Patent Nos. 6,020,196 and 5,744,360. Furthermore, tissue may be isolated from any suitable location on an animal, depending upon the type of tissue being isolated. For example, adipose tissue may be isolated from locations including, but not limited to, the tail head, the omentum or other abdominal location, subcutaneously, the stomach, hips or thighs. As used herein, the tail head region is the general area from the midline lateral and cranial to the insertion of the tail into the body of the animal, extending forward to the area of the loin and the points of the hips. Umbilical cord matrix is typically isolated from the matrix within the umbilical cord, otherwise referred to as Wharton's jelly.

Of particular interest is the use of adipose tissue obtained during surgical procedures, and especially procedures that relate to the spaying and neutering of animals such as dogs and cats. Adipose tissue collected from young dogs and cats undergoing spaying and neutering will produce a higher level of stem cells. Such cells can be used, e.g., for "banking" the cells for

subsequent return to the patient or in the allogeneic treatment of another animal.

A tissue is processed to release cells from other tissue components by any of a variety of different means or combinations thereof. In many embodiments, tissue is physically processed, *e.g.*, by cutting or mincing a tissue sample into smaller pieces. In certain embodiments, tissue is processed by exposure to an enzyme preparation that facilitates the release of cells from other tissue components, while in other embodiments, the processing of tissue does not include exposure to an enzyme that facilitates the release of cells from other tissue components. In one embodiment, the enzyme preparation is a collagenase preparation or comprises collagenase. In related embodiments, the enzyme preparation comprises one or more of trypsin-like, pepsin-like, clostripain, and neutral protease-type enzymes. Typically, the methods of the invention include processing by one or more of the following procedures:

physical cutting, enzymatic treatment, ultrasonic energy treatment, and perfluorocarbon treatment.

In one embodiment, the processing of a tissue comprises physically cutting the tissue into smaller pieces. Cutting may be performed by any means available, including, *e.g.*, the use of scissors, scalpels, razor blades, needles, and other sharp instruments.

In one embodiment, physical cutting of a sample is performed using a device comprising an array of screens. A tissue sample is placed onto the array of screens, which are typically held within a container. The number of screens arrayed together to receive the adipose tissue is preferably sufficient to contain the volume of the tissue sample within the openings of the screens. In one embodiment, the screens comprise a sharp edge, which cuts the tissue sample as it is applied to the array of screens. In another embodiment, the device is constructed such that one or more of the screens can be separated from a neighboring screen, and a cutting device, such as a wire, blade, or scalpel may be passed or inserted between the screens to further process and

cut the tissue. The screens themselves may be constructed from any suitable material, including, *e.g.*, metal, ceramic, plastic or glass. In various embodiments, the container holding the screens comprises an opening through which the tissue sample may be applied to the screens.

5 Without wishing to be bound to any particular theory, it is understood that the tissue, *e.g.*, adipose tissue, is dispersed on the openings of the screens and, therefore, has a more uniform particle distribution compared to the traditional use of scissors to mince the tissue into smaller particles. This method also avoids subjecting the tissue to high shear forces in order to
10 produce smaller particles. The adipose tissue thus becomes dispersed into more uniformly sized particles that will be enzymatically modified at a uniform rate upon subsequent or concurrent enzymatic treatment. Such arrangement promotes a more rapid release of cells and, therefore, reduces the contact time between released cells and the enzyme solution. Moreover, using the inventive
15 device and process, it also is possible to remove the free cells, since the residual tissue matrix is retained on the screens. Still further, and in contrast to heretofore known devices, it is possible to visually inspect the amount of tissue remaining on the screens, in order to assess the extent of the dissolution process and determine the point at which processing is completed.
20 Consequently, this screen device and methods utilizing the device allow for more expeditious processing of tissue, since variability of the enzyme solution does not need to be determined before hand due to the ability to see the extent of the processing by observing the amount of tissue remaining on the screens.

 In certain embodiments, processing of the tissue includes
25 enzymatic treatment, as described, *e.g.*, in Example 1. Typically, such enzymatic treatment involves exposing the tissue to one or more enzymes that facilitate the release of cells from other tissue components. Example of such enzymes include matrix metalloproteinases, clostripain, trypsin-like, pepsin-like, neutral protease-type and collagenases. Suitable proteolytic enzymes are
30 described in U.S. Patent Nos. 5,079,160; 6,589,728; 5,422,261; 5,424,208; and

5,322,790. In one embodiment, a tissue sample is exposed to collagenase at a concentration in a range of 0.01 to 10.0 mg/ml, 0.05 to 10 mg/ml, 0.5 to 2.5 mg/ml, or 0.75 to 2.0 mg/ml, for a time sufficient to release cells from other tissue components. In a related embodiment, the level of collagenase is 0.75 mg/ml (0.075%). The actual usage level may be routinely determined by the skilled artisan for the particular tissue type being digested, and it is further understood that the concentration may vary depending upon the particular source of the enzyme. In particular embodiments, collagenase is used at approximately 0.75 or 0.9 mg/ml (Sigma-Aldrich, Cat. # 2674), or 0.75 or 2.0 mg/ml (Serva NB4). Enzymatic treatment may be performed at a variety of different temperatures and time durations, which are understood generally to be inversely correlated to some degree. For example, in one embodiment, collagenase treatment is performed at 37°C for 2-5 minutes multiple times (with removal of cells after each time period) or as long as 3-4 hours. In one embodiment, the total incubation with enzyme is 20-60 minutes.

In one embodiment, ultrasonic energy is used to process a tissue sample. In a specific embodiment, a transducer is applied to a fluid filled chamber containing the tissue being processed. The energy is applied and dissolution of the tissue occurs. In related embodiment, this procedure is performed separately or in combination with enzymatic treatment. Conditions of the ultrasonic treatment are selected so that adipose tissue is affected without the cells therein being significantly damaged. The use of ultrasonic energy has previously been shown to improve the dissolution of adipose tissue under *in vivo* procedures relating to lipoaspiration and suitable conditions for *in vivo* dissolution of adipose tissue have been described in US Patent Application Publication No. 2002/0128592 A1, which conditions may be adapted for the *in vitro* uses described herein.

In another embodiment, processing of a tissue sample includes treatment with a medically-compatible perfluorocarbon solution, *e.g.*, as described in Example 2. Typically, the adipose tissue is placed into contact

with or mixed with the perfluorocarbon solution for sufficient time to generate an emulsion. The perfluorocarbon solution layer is then aspirated, leaving the aqueous layer containing the stem cells. The use of medically-compatible compositions of perfluorocarbons has been reported to aid in the *in vivo* removal
5 of adipose tissue performed on human subjects (see, *e.g.*, U.S. Patent No. 6,302,863), and methods and perfluorocarbon solutions described therein may be applied to the *in vitro* methods of the present invention.

In various embodiments, released cells are purified from other tissue components after or concurrent with the processing of a tissue sample.
10 As used herein, purification of cells means the release of cells from their normal tissue environment and does not indicate that the cells are purified or isolated from all other tissue components. In certain embodiments, purification of cells comprises separating cells from certain insoluble tissue components, including residual tissue material, such as lipids. Cells are separated from other tissue
15 components by any means known and available in the art, including, *e.g.*, the use of density gradients, centrifugation, and filtration or combinations thereof. Example of specific methods of purifying cells are known and described in the art, *e.g.*, in U.S. Patent No. 6,777,231. In certain embodiments, negative separation methods are employed to remove one or more particular types of
20 cells.

Cells prepared according to the methods of the invention may be used immediately or stored prior to use. In certain embodiments, cells are isolated from a tissue sample at a geographic location different from the location where the tissue sample was obtained or where the tissue sample is to
25 be provided to a patient. In such circumstances, the purified cells are typically stored prior to shipment to a physician or veterinarian for administration to a patient. The cells may be stored temporarily at approximately 4°C, or the cells may be frozen under liquid nitrogen for long term storage. A variety of methods of freezing cells for long term storage and recovery are known in the art and
30 may be used according to the invention, including freezing cells in a medium

comprising fetal bovine serum and dimethylsulfoxide (DMSO), as exemplified in Example 7.

In certain embodiments, purified cells, whether previously frozen or not, are placed into a vehicle suitable for administration. For example,
5 purified cells may be placed into a syringe suitable for injection into a patient at a wound site or via intravenous administration.

Remarkably, it was discovered according to the invention that purified cell populations retained viability over time, when stored under refrigeration, *e.g.*, at temperatures less than 12°, or while shipped on cold
10 packs. This discovery was particularly surprising, since it had previously been shown that the use of enzymatic treatments like collagenase reduce the viability of cells, due to degradation of membrane structures (Brundstedt, J. *et al.*, Methods in Diabetes Research, V. 1, Laboratory Methods, Larner, J. and Pohl, S.L. (eds.), Wiley-Interscience, New York, 1985), and the prior understanding in
15 the art was that purified stem cell populations could not be stored without substantial loss of viability unless specific steps were taken to preserve viability, including addition of proteins, nutrients, serum and tissue culture media of a variety of compositions.

Accordingly, in another embodiment, the invention provides a
20 purified cell population comprising stem cells useful in the treatment and prevention of injury and disease, which can be stored at a temperature of less than 12°C and transported under refrigeration or in the presence of cold packs prior to delivery to a patient. Such stem cell populations and compositions may be used in procedures providing purified stem cells for therapeutic or
25 prophylactic purposes.

In certain embodiments, such cell populations are prepared according to a method of the present invention. However, in other embodiments, the cell populations may be isolated by any other means in the art, including those methods that employ enzymatic treatment and those that do
30 not. In one embodiment, the cell populations and related methods include a

step of culturing or rinsing the cells in the presence of serum or nutrient buffers, which can inhibit collagenase activity due to serum components.

The invention also includes methods of providing a stem cell population that may be transported under refrigeration or on cold packs to a location other than where the stem cell population was prepared. In one embodiment of such a procedure, a tissue sample is obtained from a patient by a physician or veterinarian and shipped to a laboratory. At the laboratory, the sample is processed to provide a purified cell population comprising stem cells. The purified cell population, or a portion thereof, is then shipped to a physician or veterinarian and subsequently delivered to the patient. The purified cells may be stored prior to shipment, *e.g.*, at 4°C or under liquid nitrogen. In one embodiment, the purified cell population is transferred to and shipped in a vehicle, such as a syringe, suitable for delivering the cell population to a patient. In particular embodiments, the cells are stored cold, *i.e.* less than 12°C for 24 hours or 48 hours without losing significant viability or losing less than 5%, 10%, 20%, or 50% viability.

The current invention may be further applied industrially as a method of providing a medical service. Thus, in one embodiment, the invention includes a method of providing purified stem cells, comprising providing a kit for obtaining a tissue sample from a patient to a physician or veterinarian, processing the obtained tissue sample to purify a cell population comprising stem cells, and shipping a purified cell population to a physician or veterinarian in a device suitable for administering the cell population to a patient. These methods permit such off-site processing, since they provide a stem cell population having a high percentage of viable cells even after storage and shipment at less than 12°C or on ice. Of course, it is understood that the method may be modified without falling without the scope of the present invention. For example, in one embodiment, the method may not include the step of providing a kit for obtaining a tissue sample, since the physician or veterinarian can usually obtain and ship a tissue sample to a laboratory for

processing using materials at hand. Furthermore, it is not necessary according to the method for the cells to be shipped in a device suitable for administration to a patient. Rather, the cells may be shipped in a container, *e.g.*, a vial, and subsequently transferred into a device suitable for administration.

5 In one specific embodiment, a sample of adipose tissue obtained from a horse by a veterinarian is processed according to a method of the invention, and the purified stem cell population is placed into a syringe and shipped on cold packs to the veterinarian. The purified stem cell population may then be provided to a site of injury or potential injury in the horse from
10 which the tissue sample was obtained. In particular embodiments, the site of injury is a tendon, ligament, cartilage or bone, including, *e.g.*, a strain or fracture.

B. Compositions and Kits Comprising Purified Stem Cells

 The method of the invention results in a purified cell population
15 having a unique composition that has distinct advantages in the treatment and prevention of injuries. The cell population purified according to the methods of the invention includes multipotent stem cells, such as *e.g.*, mesenchymal or embryonic stem cells. However, as used herein, the term “purified” does not indicate the presence of only stem cells. Rather, the term “purified” indicates
20 that the cells are removed from their natural tissue environment and are present at a higher concentration as compared to the normal tissue environment. Accordingly, a “purified” cell population may further include cell types in addition to stem cells and may include additional tissue components. In particular embodiments, purified cell populations comprise at least 10,000, 20,000,
25 50,000, 100,000, 200,000, 500,000, 600,000, 700,000, 800,000, 900,000, 1×10^6 , 2×10^6 , 3×10^6 , 5×10^6 , or 10×10^6 cells per gram of tissue. In certain embodiments, at least 600,000 to 70×10^6 cells are isolated from 3 to 50 grams of tissue. In related embodiments, the purified cells are present at a

concentration approximately 40 to 50-fold greater than their concentration in the tissue from which they were isolated, when initially pelleted.

In certain embodiments, a stem cell is of mesodermal origin. Typically, such stem cells retain two or more mesodermal or mesenchymal developmental phenotypes. In particular, such cells have the capacity to develop into mesodermal tissues, such as mature adipose tissue, bone, various tissues of the heart, dermal connective tissue, hemangial tissues, muscle tissues, urogenital tissues, pleural and peritoneal tissues, viscera, mesodermal glandular tissue and stromal tissue. In other embodiment, a stem cell has the capacity to develop into neural ectodermal tissue.

The purified cells demonstrate a high degree of viability, both before and after storage at 4°C or under liquid nitrogen, and after being shipped at temperatures less than 12 °C or on ice. In certain embodiments, the percentage of viable cells, as determined by standard Trypan blue dye exclusion methods, immediately following preparation of a tissue sample is at least 50%, 60%, 70%, 80% or 90%. In related embodiments, the percentage of viable cells, as determined by standard Trypan blue dye exclusion methods, following storage at 4°C for 24 hours or storage under liquid nitrogen for two weeks is at least 40%, 50%, 60%, 70%, 80% or 90%. In another embodiment, the percentage of viable cells, as determined by standard Trypan blue dye exclusion methods, following refrigerated storage or shipment on ice packs for less than 24 hours is at least 30%, 40%, 50%, 60%, 70%, 80% or 90%.

In addition, in certain embodiments, the cell populations also include other cell types, such as one or more of the following: red blood cells, white blood cells, neutrophils, monocyte/macrophages, fibroblasts, fibroblast-like cells, lymphocytes, and basophils. However, in certain embodiments, the compositions and cell populations do not include lymphocytes (*i.e.*, T or B cells) or have a significantly reduced percentage of lymphocytes as compared to the amount present in peripheral blood. In specific embodiments, the percent of total cells in the purified cell population that are lymphocytes is reduced by at

least 50%, 60%, 70%, 80%, 90%, 95%, 99% or 100% as compared to the percent of total cells in the original tissue sample that are lymphocytes. In related embodiments, lymphocytes represent less than 1%, 2%, 5%, 10%, 20%, 30%, 40%, or 50% of the total cells present in the purified cell population.

5 In particular embodiments, the purified cell population does not comprise an appreciable number of lymphocytes. An appreciable number of lymphocytes, as used herein, refers to at least 5% of the cell population being lymphocytes. Since the methods of the invention do not typically include a step of separating stem cells from other purified cells, these additional cells may be present in the
10 originally purified cell population. Alternatively, non-stem cells may be added to the purified cell population at any time prior to administration to a patient.

In further embodiments, the cell populations also include non-cellular tissue components. Such non-cellular components may be soluble factors, or, alternatively, they may be insoluble components, such as lipids, or
15 both. Examples of such non-cellular tissue components include extracellular matrix proteins, proteoglycans, secreted factors, cytokines, growth factors, differentiation-inducing factors, and differentiation-inhibiting factors, or fragments thereof. In one embodiment, the cell populations include collagen, thrombospondin, fibronectin, vitronectin, laminin, or fragments thereof. In a
20 particular embodiment, the cell populations include collagen or fragments thereof. Collagens include, but are not limited to, Type I, Type II, Type III, and Type IV collagen. Again, these additional non-cell components frequently will be present in the originally isolated cell population. However, in certain
25 embodiments, such non-cell components are added to the purified cell population prior to administration to a patient.

Without wishing to be bound to any particular theory, it is understood that the presence of tissue components in addition to stem cells provides a therapeutic advantage over stem cell populations lacking other tissue components, *e.g.*, by providing additional factors that promote
30 appropriate differentiation of the stem cells upon administration to a patient

and/or possess. In addition, certain components are understood to possess intrinsic wound healing and preventative properties and, thus, cooperate with the stem cells in tissue repair and wound prevention.

In certain embodiments, the purified cell populations are present
5 within a composition adapted for and suitable for delivery to a patient, *i.e.*, physiologically compatible. Accordingly, compositions of the stem cell populations will often further comprise one or more buffers (*e.g.*, neutral buffered saline or phosphate buffered saline), carbohydrates (*e.g.*, glucose, mannose, sucrose or dextrans), mannitol, proteins, polypeptides or amino acids
10 such as glycine, antioxidants, bacteriostats, chelating agents such as EDTA or glutathione, adjuvants (*e.g.*, aluminum hydroxide), solutes that render the formulation isotonic, hypotonic or weakly hypertonic with the blood of a recipient, suspending agents, thickening agents and/or preservatives.

In other embodiments, the purified cell populations are present
15 within a composition adapted for or suitable for freezing or storage, such as the freezing medium described in Example 7.

The methods and compositions of the present invention are particularly well-adapted to being practiced using a kit, since they permit the storage and shipment of stem cell populations. In certain embodiments, a kit
20 comprises a device suitable for administering the purified stem cell composition to a patient and containing an amount of stem cell composition to be administered. In one embodiment, a kit useful in the treatment of a musculo-skeletal tissue injury in an animal comprises a syringe containing a composition comprising purified adipose tissue-derived stem cells obtained from the animal
25 in a physiologically compatible solution. It is understood that a kit may include any of the purified stem cell populations and compositions described herein. Accordingly, kits of the invention may be prepared for autologous, allogeneic or xenogeneic administration, and may further comprise additional tissue components (cellular or non-cellular) that are co-purified with the stem cells or
30 added to the composition after purification of the stem cells.

C. Methods of Treating and Preventing Injury

In certain embodiment, the purified stem cells and compositions comprising the same are used to treat a clinically obvious injury or disease in a patient. In other embodiment, they are used prophylactically to prevent sub-clinically non-obvious injury or disease. In addition, in certain embodiments, they are used autologously to treat a patient from which the purified stem cells were isolated, while in other embodiments, they are used allogeneically to treat a patient other than the donor from which the stem cells were purified. In one embodiment, they are used to treat a patient of the same species, while in another embodiment, they are used to treat a patient of a difference species, *i.e.*, xenogeneic.

In certain embodiments, the purified stem cells and related compositions are used to treat a variety of different diseases, including but not limited to inflammatory diseases, cardiovascular diseases, nervous system diseases, tumors, demyelinating diseases, digestive system diseases, endocrine system diseases, reproductive system diseases, hemic and lymphatic diseases, immunological diseases, mental disorders, musculoskeletal diseases, neuromuscular diseases, metabolic diseases, skin and connective tissue diseases, urological diseases.

In various embodiments, the purified stem cells and related compositions are used to treat a variety of different wounds, including but not limited to, abrasions, avulsions, blowing wounds, incised wounds, burns, contusions, puncture wounds, surgical wounds and subcutaneous wounds.

In particular embodiments, the purified stem cells and related compositions are used to treat or prevent a variety of injuries, including but not limited to, injuries to muscle, connective tissue (including tendon, ligament and cartilage), bone, hoof laminae, lung tissue, blood vessels, nerve, liver, musculoskeletal tissue or cardiac tissue. In particular embodiments, the injury is a sports related injury, which includes but is not limited to contusions, myositis, strains, (including muscle and tendon strains), microtears, fractures (including

avulsion fractures), dislocation, tear, sprains, stress fractures, bursitis, and articular cartilage injury.

In one embodiment, the injury is an injury associated with a performance animal, such as a tendon or ligament injury, which are frequently observed in competitive or racing mammals such as humans, horses, dogs and camels. In one embodiment, the injury occurs in a horse or camel within a superficial digital flexor tendon, suspensory ligament, accessory ligament of the deep digital flexor tendon, menisci, or other ligament such as cruciate ligaments. In another embodiment, the injury occurs in a dog within an Achilles tendon, cruciate ligament, meniscus, or flexor tendon. In certain embodiments wherein the patient is a human, the injury is of the Achilles tendon, quadriceps tendon, rotator cuff, lateral or medial epicondylitis, cruciate ligament, intervertebral disc or meniscus. Accordingly, in related embodiments, examples of particular injuries include tendonitis, tendinopathy, desmitis, bowed tendon, fractures, and strains.

Cell populations and related compositions may be provided to a patient by a variety of different means. In certain embodiments, they are provided locally, *e.g.*, to a site of actual or potential injury. In one embodiment, they are provided using a syringe to inject the compositions at a site of possible or actual injury or disease. In other embodiments, they are provided systemically. In one embodiment, they are administered to the bloodstream intravenously or intra-arterially. The particular route of administration will depend, in large part, upon the location and nature of the disease or injury being treated or prevented. Accordingly, the invention includes providing a cell population or composition of the invention via any known and available method or route, including but not limited to oral, parenteral, intravenous, intra-arterial, intranasal, and intramuscular administration.

In one specific embodiment, a method of treatment comprises injecting a composition comprising stem cells isolated from an adipose tissue sample obtained from the tail head region of a horse and prepared according to

a method of the invention into the same horse at a site of actual or potential injury, such as a tendon or ligament.

The development of suitable dosing and treatment regimens for using the cell populations and compositions described herein in a variety of treatment regimens, including *e.g.*, oral, parenteral, intravenous, intranasal, and intramuscular administration and formulation, will again be driven in large part by the disease or injury being treated or prevented and the route of administration. The determination of suitable dosages and treatment regimens may be readily accomplished based upon information generally known in the art.

Treatment may comprise a single treatment or multiple treatments. In particular, for preventative purposes, it is contemplated in certain embodiments that purified cell populations of the invention are administered prior to a stress that might potentially cause injury, such as, *e.g.*, an animal race (*e.g.*, dog or horse race).

All of the above U.S. patents, U.S. patent application publications, U.S. patent applications, foreign patents, foreign patent applications and non-patent publications referred to in this specification and/or listed in the Application Data Sheet, are incorporated herein by reference, in their entirety.

EXAMPLE 1

PREPARATION OF ADIPOSE-TISSUE DERIVED STEM CELLS USING ENZYMATIC DEGRADATION

Stem cells from adipose tissue obtained from four equines were prepared according to the following procedure, and the number and viability of purified cells was determined.

- a. Label processing tubes (50 ml conicals). Determine the weight of each conical, record on the label.
- b. Transfer fat sample to the inside surface of a lid from a Petri dish. Allow to drain.

- c. Place fat sample in pre-weighed conical tube.
- d. Record weight of fat sample and tube.
- e. Add 30-40 ml of sterile PBS to the conical, cap and gently invert several times.
- 5 f. Carefully pour off liquid by using sterile forceps to retain the fat.
- g. Transfer fat to the bottom portion of a sterile petri dish and mince with scissors and forceps.
- h. Return minced sample to original 50 ml conical by using
10 sterile scoop.
- i. Rinse the petri dish with 15 ml of sterile PBS to dislodge any remaining fat particles and pour liquid/particles into the conical.
- j. Carefully aspirate PBS from the tube until approximately 5 ml of PBS remains.
- 15 k. Prepare the required amount of collagenase solution by determining the volume of the fat present in the conical. Make sufficient volume of collagenase to equal 1.1 X the volume of fat to be treated. Collagenase stock is 0.075% in PBS (*i.e.*, 75 mg per 100 ml). Filter sterilize the collagenase stock solution by using a 0.22 μ m sterile filter. Place collagenase stock solution in a
20 37° C water bath.
- l. Add enough sterile PBS to bring level to 40-45 ml mark, then cap.
- m. Mix by inverting several times.
- n. Aspirate the aqueous layer present, leaving 5 ml of volume.
- 25 o. Repeat Steps 2l-2n until aqueous layer is relatively clear of debris/blood (*e.g.*, 2x or 3x, depending on how bloody the sample is).
- p. After final aspiration, add collagenase enzyme solution (warmed to 37 °C). Mix well. Add a volume of collagenase stock solution equal to the total volume of the minced fat sample (fat and residual PBS).

- q. Incubate tubes in 37 °C water bath with agitation for 20-60 minutes, depending on the rate of fat degradation.
- r. Dry tubes and spray outside of tubes with 70% Isopropyl alcohol.
- 5 s. Add enzyme-neutralizing solution (optional, based on adding DMEM with FCS).
- t. Centrifuge tube at 400 x g for 5 minutes, swinging bucket rotor (set at 4 °C).
- u. Aspirate/pour off supernatant, removing fat layer.
- 10 v. Resuspend pellet by gently "flicking" the bottom of the tube with a finger.
- w. Add 10 ml of PBS and mix gently by inversion.
- x. Pour resuspended cells through tissue filter (*e.g.*, 70 µm) into a labeled 50 ml conical tube to remove any tissue matrix and debris. After
- 15 the suspension has drained into the conical, gently rinse the tissue filter with approximately 20 ml of sterile PBS, collecting the rinsate in the conical (including the underside of the filter unit itself).
- y. Centrifuge the tube at 400 x g for 5 minutes, swinging bucket rotor.
- 20 z. Aspirate/pour off supernatant.
- aa. Resuspend cell pellet in a small volume of medium.
- bb. Transfer 50 µl of each sample to the appropriate labeled 0.5 ml plastic tube for cell counting and viability determination.
- Cell count and viability determinations were performed according
- 25 to the following procedures.
- a. Add 50 µl of 0.4% Trypan Blue dye exclusion medium to the tube in Step bb.
- b. Mix gently and allow to stand for 1-2 minutes.
- c. Load one chamber of the hemacytometer with a sample.
- 30 Let sit for 1-2 minutes.

d. Perform cell count and viability determination by counting at least 100 cells (but fewer than 500 cells) contained in the 4 large grid areas.

e. Perform calculations of cell number and viability.

The results of these experiments are shown in Table 1, which provides the number of cells isolated per gram of adipose tissue and the percent viability of cells from each sample.

Table 1. Viability and number of adipose tissue-derived cells

Sample Number	% Viability	Cell Number/g of adipose tissue
1	68.5	0.86×10^6
2	55.7	0.91×10^6
3	72.0	0.36×10^6
4	75.4	0.35×10^6

These experiments demonstrate that the procedure of the present invention provides a remarkably high number of viable cells per gram of tissue and is, therefore, extremely useful for preparing stem cells for therapeutic and prophylactic treatments.

EXAMPLE 2

PREPARATION OF ADIPOSE-TISSUE DERIVED STEM CELLS USING PERFLUOROCARBONS TO RELEASE STEM CELLS FROM TISSUE

The following procedure for preparing adipose-tissue derived stem cells is adapted from the procedure of Example 1 by the addition of treatment with perfluorocarbons to assist in the release of stem cells from adipose tissue.

- a. Label processing tubes (50 ml conicals). Determine the weight of each conical, record on the label.
- b. Transfer fat sample to the inside surface of a lid from a Petri dish. Allow to drain.
- c. Place fat sample in pre-weighed conical tube.
- d. Record weight of fat sample and tube.

- e. Add 30-40 ml of sterile PBS (with a pH of 6.9; to be used throughout the procedure) to the conical, cap and gently invert several times.
- f. Carefully pour off liquid by using sterile forceps to retain the fat.
- 5 g. Transfer fat to the bottom portion of a sterile Petri dish and mince with scissors and forceps.
- h. Return minced sample to original 50 ml conical by using sterile scoop.
- i. Rinse the Petri dish with 15 ml of sterile PBS to dislodge
10 any remaining fat particles and pour liquid/particles into the conical.
- j. Carefully aspirate PBS from the tube until approximately 5 ml of PBS remains.
- k. Add an amount of perfluorocarbon solution equal to the volume of the adipose sample, including PBS. Shake to bring the adipose
15 tissue into contact with the perfluorocarbon solution. Use of a "rocker" platform will aid in this procedure. At completion, aspirate the PFC layer without removing the aqueous layer.
- l. Prepare the required amount of collagenase solution by determining the volume of the fat present in the conical. Make sufficient volume
20 of collagenase to equal 1.1 X the volume of fat to be treated. Collagenase stock is 0.075% in PBS (i.e., 75 mg per 100 ml). Filter sterilize the collagenase stock solution by using a 0.22 μ m sterile filter. Place collagenase stock solution in a 28 °C water bath.
- m. Add enough sterile PBS to bring level to 40-45 ml mark,
25 then cap.
- n. Mix by inverting several times.
- o. Aspirate aqueous layer, leaving 5 ml of volume .
- p. Repeat Steps 2l-2n until aqueous layer is relatively clear of debris/blood (e.g., 2x or 3x, depending on how bloody the sample is).

- q. After final aspiration, add collagenase enzyme solution (warmed to 37 °C). Mix well. Add a volume of collagenase stock solution equal to the total volume of the minced fat sample (fat and residual PBS).
- 5 r. Incubate tubes in 37 °C water bath with agitation for 20-60 minutes, depending on the rate of fat degradation.
- s. Dry tubes and spray outside of tubes with 70% Isopropyl alcohol.
- t. Add enzyme-neutralizing solution (optional, based on adding DMEM with FCS).
- 10 u. Centrifuge tube at 400 x g for 5 minutes, swinging bucket rotor (set at 4 °C).
- v. Aspirate/pour off supernatant, removing fat layer.
- w. Resuspend pellet by gently "flicking" the bottom of the tube with a finger.
- 15 x. Add 10 ml of PBS and mix gently by inversion.
- y. Pour resuspended cells through tissue filter (70 µm) into a labeled 50 ml conical tube to remove any tissue matrix and debris. After the suspension has drained into the conical tube, gently rinse the tissue filter with approximately 20 ml of sterile PBS, collecting the rinsate in the conical
- 20 (including the underside of the filter unit itself).
- z. Centrifuge the tube at 400 x g for 5 minutes, swinging bucket rotor.
- aa. Aspirate/pour off supernatant.
- bb. Resuspend cell pellet in a small volume of medium.
- 25 cc. Transfer 50 µl of each sample to the appropriate labeled 0.5 ml plastic tube for cell counting and viability determination.

EXAMPLE 3ISOLATION OF STEM CELLS FROM ADIPOSE TISSUE OBTAINED FROM THE TAIL HEAD
REGION OF A HORSE

The optimal location on a horse for the collection of an adipose
 5 tissue sample for isolation of stem cells is not readily apparent, since fat is
 deposited at numerous sites in horses. The system of body condition scoring
 (Ott EA. Chairman Subcommittee on horse nutrition: Nutritional Requirement of
 Horses. 5th ed National Academy Press, Washington DC (1989)) describes fat
 accumulating in numerous locations on the horse.

10 To determine if the tail head region is a preferred location for
 collecting fat tissue, a number of adipose tissue samples were obtained from
 both thin and fat horses and processed as described in Example 1.
 Specifically, samples were obtained from the tail head region of four equine
 patients and the neck of one equine patient. The processed samples were
 15 analyzed to determine viability and number of stem cells, and the results are
 provided in Table 2. Adherent cells were observed for selected samples that
 were cultured, indicating the presence of viable stem cells.

Table 2. Viability and Number of Cells Obtained from the Tail Head Region

Sample #	Location	% Viability	Cell Number/g of adipose tissue
20 1	Tail Head	68.5	0.86×10^6
2	Tail Head	55.7	0.91×10^6
3	Tail Head	72.0	0.36×10^6
4	Tail Head	75.4	0.35×10^6
5	Neck	N/A	No cells isolated

25 The results of these experiments establish that the tail head
 region is a preferred location to obtain adipose tissue from a horse. Adipose
 tissue derived from the tail head region provided increased number of cells as
 compared to adipose tissue derived from the neck. The tail head region also
 offers additional advantages as compared to other locations, including greater
 30 ease of access and safety in a chute or stanchion, no requirement for general

anesthesia, no significant nerves or vessels are located in this area, and fat is available in adequate amounts for collection and is close to the body surface. In addition, the surprising finding was made that the tail head area is one of the few areas where fat is located distinct from other fibrous tissues.

5

EXAMPLE 4

Small animals, including dogs and cats, possess a number of potential anatomical sites for obtaining adipose tissue. Stem cells were isolated from adipose tissue obtained from a variety of anatomical sites in animals according to the procedure described in Example 1, and the amount of cells isolated and their viability was determined. The results are shown in Table 3.

Table 3. Viability and Number of Cells Obtained from Small Animals

Identification No.	Species	Sample Type	% Viability	Viable Cells/gram Adipose Tissue
061302-01	Feline	Lipoaspirate	85.0	N/A
061302-02	Canine	Lipoaspirate	77.0	N/A
061902-01	Feline	Lipectomy	89.4	2.85×10^6 (total)
071202-01	Canine	Lipectomy	97.1	1.12×10^6 (total)
110702-01	Feline	Lipectomy	94.4	0.61×10^6
092303-01	Canine	Broad-Lig.	86.5	2.3×10^6
092303-02	Canine	Subcutaneous	75.2	0.59×10^6
092303-03	Canine	Omental	55.2	1.1×10^6
092403-01	Feline	Subcutaneous	63.2	0.26×10^6
092403-04	Feline	Omental	66.7	0.30×10^6
092403-02	Canine	Omental	85.2	1.84×10^6
092403-03	Feline	Subcutaneous	91.7	0.8×10^6
092503-01	Feline	Subcutaneous	80.6	0.26×10^6
092503-02	Feline	Omental	77.0	0.51×10^6
092603-01	Canine	Subcutaneous	76.1	0.46×10^6
092603-02	Canine	Omental	90.3	1.57×10^6
100303-01	Canine	Subcutaneous	68.6	N/A
100303-02	Canine	Omental	67.0	0.21×10^6
100303-03	Canine	Broad Lig.	60.0	0.31×10^6
100303-04	Canine	Broad Lig.	67.0	0.22×10^6

These results demonstrate that viable stem cells may be isolated from a variety of different locations in small animals. In addition, they further establish that the procedures of the present invention result in the isolation of a surprisingly

15

substantial number of viable cells and are, therefore, particularly well-suited for the preparation of stem cells for therapeutic and prophylactic treatment of patients, since a large number of viable cells may be isolated from a relatively small tissue sample.

5

EXAMPLE 5

PREPARATION OF ADIPOSE TISSUE-DERIVED STEM CELLS USING A MODULE FOR TISSUE PROCESSING

Adipose tissue obtained from a patient by standard methods (*e.g.*,
10 lipectomy, lipoaspiration, or other suitable procedure) is brought into contact with a series of screens within a module and is forced into the mesh of the screens. Once the adipose tissue has been dispersed throughout the screens, the individual screens are separated slightly from their neighbors, creating a gap between the screens. The screens are moved in a parallel manner relative
15 to one another, or a thin blade may be passed between adjacent screens in order to obtain discretely distributed adipose tissue held within the openings of each screen.

The set of screens is processed individually or is maintained in a “block” arrangement. In either case, an enzymatic solution is added to the
20 container in which the screens are placed and allowed to come into contact with the adherent adipose tissue. Free cells released from the screens are recovered, while the residual tissue matrix is retained on the screens.

EXAMPLE 6

ISOLATION OF STEM CELLS USING ADHERENT MATERIAL

25 Stem cells were isolated from adipose tissue using small adherent materials, *e.g.*, packing peanuts or Velcro pieces, to which the adipose tissue adheres, according to the following procedure. The effect of including either of

these adherent materials in samples being processed was determined by monitoring their effect on filtration rates, as described below.

a. Packing peanuts were "chopped" with a vegetable chopper and scissors to reduce their nominal size to between 1 mm to 10 mm in diameter approximately.

b. The chopped peanuts were placed in a beaker and water was added. Only those peanuts that appeared to be within the dimensional range and floated were recovered.

c. Velcro was cut into pieces, essentially creating shreds of Velcro approximately 2-4 mm wide and the length of the original Velcro strip width. These shreds were placed in water to see if they floated.

d. The chopped peanuts (approx. 1 gram) were placed in the top of a Corning 115 mL filtration system and 100 mL of water was added. Vacuum was applied and the time that it took to filter the 100 mL was determined. As a control, the time to filter 100 mL of water by itself was measured.

e. In a fresh filtration device, the time to filter 100 mL of water in the presence of the shredded Velcro (approx. 1 gram) was determined.

f. 20 g of fat was minced as described in Example 1. The minced tissue was placed in a fresh Corning 115 mL filtration device. 100 mL of water was added, the fat particles were dispersed. Vacuum was applied and the time to filter was determined.

g. 20 g of fat was minced as described in Example 1. The minced tissue was mixed with the chopped peanuts and placed in a fresh Corning 115 mL filtration device. 100 mL of water was added and the particles were dispersed. Vacuum was applied and the time to filter the 100 mL of water was determined.

h. 20 g of fat was minced per the standard protocol. The shredded Velcro pieces were mixed with the fat and placed in a fresh Corning 115 mL filtration system. 100 mL of water was added and the particles were

dispersed. Vacuum was applied and the time to filter the 100 mL of water was determined.

Most of the Velcro pieces and all of the minced packing peanut pieces floated in water. The time for 100 mL of water to filter in a filter system was recorded, along with the time it took for 100 mL of water to filter in the same filter unit in the presence of either the Velcro or packing peanut pieces. The final filtration assessment was determined with the fat tissue added directly to the filter or added to the filter after being mixed with the Velcro or packing peanut pieces. The fat coated both the Velcro and the packing peanuts quite well without extensive mixing being required, and it appeared that nearly all of the minced fat was in contact with the Velcro or packing peanut pieces. The initial filtration results are shown in Table 4.

Table 4. Filtration Times of Adipose Tissue Samples in the Presence or Absence of Adherent Materials

Filter Unit	Condition	Time to Filter (sec)
1	H ₂ O	25
	H ₂ O/Velcro	25
	H ₂ O/Velcro/Fat (20.361 g)	39
2	H ₂ O	24
	H ₂ O/Packing Peanuts	25
	H ₂ O/Packing Peanuts/Fat (20.545 g)	37
3	H ₂ O	27
	H ₂ O/Fat (20.605 g)	53

Additional filtration studies were performed using a 0.2 µm filter system from Nalgene. The previous assessment was made with a Corning filter system. Approximately 1 g of Velcro pieces and polystyrene packing peanut pieces were used in each of three filter units tested per condition. Fat without any agent present was used as the control. The time for 100 mL of water to filter in a filter system was assessed in the absence (control) or presence of either the Velcro/fat or packing peanut pieces/fat mixtures. The

time for 100 mL of water to filter in a filter system was assessed in the absence (control) or presence of either the Velcro/fat or packing peanut pieces/fat mixtures. The time for 100 mL of water to filter was determined for each filter, after which the test condition was evaluated in the same filter. Approximately 20 g of fat was used per condition. The results are shown below in Table 5.

Table 5. Filtration Times of Adipose Tissue Samples in the Presence or Absence of Adherent Materials

Filter Unit Number	Agent	Mass of Fat (g)	Time-to-Filter (sec)	
			Water Only	Water/Fat/Agent
1	None	20.156	24	44
2	None	20.176	23	48
3	None	20.511	23	51
4	Velcro	20.219	22	29
5	Velcro	20.373	22	30
6	Velcro	20.392	21	29
7	Polystrene Peanuts	20.051	23	40
8	Polystrene Peanuts	20.539	23	39
9	Polystrene Peanuts	20.534	20	34

The results shown above clearly demonstrate that the presence of either the Velcro or packing peanut pieces with the fat resulted in a substantial improvement in the filtration time. Filtration of untreated fat took almost twice as long compared to filtration of water alone, increasing from 27 seconds to 53 seconds upon the addition of the fat. However, filtration of fat mixed with the Velcro or packing peanut pieces only took approximately 1.5 times as long compared to filtration of water alone. These results indicate that the binding of the fat to the adherent material results in the generation of aggregates of adherent material/fat, which facilitates further processing.

Based on the observations made about the interaction of Velcro pieces with adipose tissue and the improved filtration that resulted from mixing the two together, an experimental assessment of the impact of including Velcro

pieces present in the tissue processing procedure of Example 1 was performed as described below.

- Approximately 1 g of Velcro pieces that had been washed with water and then dried was mixed with approximately 10.5 g of adipose tissue. A control lacking Velcro pieces was performed with 10.9 g of adipose tissue. The test materials were processed essentially according to the protocol in Example 1. Cell count and viability assays were performed at the end of the process, and the results are shown in Table 6.

Table 6. Stem Cell Viability and Cell Number When Processed Using Velcro Pieces

Condition	Viability (%)	Cells/g	Cell Yield
No Velcro	62.5	20,000	220,000
Velcro	62.6	60,000	660,000

- These results demonstrate that including small adherent materials during tissue processing results in an increased cell yield with comparable viability. Accordingly, the addition of small adherent materials during tissue processing provides a remarkably superior method for preparing stem cells, since it provides the substantial advantage of requiring less tissue to yield the same number of cells.

EXAMPLE 7

METHOD OF FREEZING PURIFIED STEM CELLS

Purified cell populations are prepared for storage in liquid nitrogen according to the following procedure.

- a. Freezing medium is prepared by determining the total number of cryovials to be used. Generally, this involves dividing the number of cells available by 3 million. Each cryovial receives 1.0 ml of freezing medium/cells, so the total number of cryovials is multiplied by 1.25 to determine the volume of freezing medium made. The appropriate number of cryovials is placed into a Cryo-Safe that has been stored at -10°C overnight.

- b. While working in the tissue culture hood, pipette the appropriate amount of DMSO and fetal bovine serum such that the final amount of DMSO in the freezing medium is 10% and the final amount of fetal bovine serum in the freezing medium is 90%. Mix well.
- 5 c. Process the freezing medium through a sterile, DMSO-compatible 0.2 μm filter into a sterile 50 ml conical tube. Place the tube in the refrigerator for at least 45 min.
- d. Centrifuge the isolated cell preparation at 400 x g for 10 min. Carefully pour off the supernatant and flick the tube to fluidize the cell
10 pellet.
- e. Add enough freezing medium to the conical tube to yield 1 ml for each cryovial. This is done by slowly adding the freezing medium over the course of 30-60 seconds, while swirling the suspension to ensure mixing.
- f. Gently pipette the cells with the freezing medium in order to
15 ensure that the cells are completely resuspended.
- g. Place 1.0 ml of the cell suspension into each of the cryovials. Cap and immediately transfer the vials to a Mr. Frosty.
- h. Immediately place the Mr. Frosty on the bottom shelf of a -80°C freezer. Place a cryovial storage cane in the freezer.
- 20 i. After a minimum of four hours storage in the -80°C freezer (not to exceed 20 hours), place the cryovials in the cane and immediately transfer the cane to a liquid nitrogen storage tank.

Cells frozen under liquid nitrogen are thawed as follows.

- a. Recovery medium is made by combining the appropriate
25 volume of autologous serum and Iscove's Modified Dulbecco's Medium (IMDM) to allow for a minimum of 1:10 dilution of the contents of the cryovial (usually 1.0 ml). To this volume, add 1.5 ml to determine the final volume of recovery medium that should be made. Preferably, the ratio is 1:15. The final recovery medium is 20% autologous serum and 80% IMDM by volume. Transfer the
30 amount of recovery medium to be used for washing to a sterile 50 ml conical

tube. Place both tubes of recovery medium in the refrigerator for a minimum of 30 min.

b. The cryovial is removed from the liquid nitrogen storage tank and immediately placed in contact with a 37°C water bath. The threads of
5 the vial should not be submerged below the surface of the water.

c. After approximately 2.5 min., the vial is examined to determine if the cells have thawed. Excess heating of the vial should be avoided.

d. As soon as the cells appear thawed, the outside of the vial
10 is washed with 70% isopropyl alcohol and placed into a Cryo-Safe that was stored at -10°C overnight.

e. The contents of the vial are immediately transferred to the 50 ml conical containing the washing recovery medium using a sterile pipette, and the suspension is gently mixed.

15 f. The cell suspension is centrifuged at 400 x g for 6 min.

g. The supernatant is gently poured off and the tube is flicked to fluidize the cells.

h. Add 1.0 ml recovery medium to resuspend the cells. Very slowly pipette the cells up and down in the pipette to mix the cells. Transfer the
20 cells to a cryovial stored in the Cryo-Safe.

i. A 30 µl aliquot of the cell suspension should be taken for determining the viability and cell count.

j. 40 µl of sterile ticarcillin stock solution (25 mg/ml) is added and mixed gently.

25 k. An 18 gauge needle attached to a sterile 1 ml syringe is used to draw the suspension into the syringe. Remove any trapped bubbles by gently tapping the barrel of the syringe.

l. Remove the needle and place the sterile syringe tip cap on the end of the syringe.

30 m. Place the syringe in the refrigerator.

Packages of cells are prepared for shipment to a physician or veterinarian as follows.

- a. A label including identification information is attached to the syringe.
- 5 b. The syringe is wrapped in bubble wrap and placed in a shipping box with ice packs above and below the syringe.
- c. Any remaining space in the box is filled with packing material, the lid is inserted, and the box is sealed.

The viability of cells frozen and thawed according to the
 10 procedure is greater than 65%, indicating that these procedures may be used successfully to store purified stem cells.

EXAMPLE 8

VALIDATION OF SYRINGE STORAGE OF CELLS

15 The viability of cells obtained by the collagenase-based processing protocol described in Example 1 was further examined to determine the impact of placing cells in a syringe and shipping the syringe to the attending veterinarian for return to the patient. The viability of each cell preparation was
 20 determined on Day 0. An aliquot of each cell preparation was then placed in a syringe, which was placed in a package with frozen cold packs and left at room temperature for between 20-24 hours before viability was determined (Day 1). For a subset of cell preparations, a control aliquot was stored in a refrigerator in
 25 a plastic tube for viability determination on Day 1. "ND" indicates that the viability was not determined. The results of these studies are provided in Table 7.

Table 7. Viability of Stem Cell Preparation Stored in a Syringe

	Viability		
	Day 0	Day 1	
Patient Identification	Control (%)	Control (%)	Syringe (%)
092403-01	63.2	ND	74.0

092403-02	85.2	ND	73.5
092403-03	91.7	ND	83.8
092403-04	66.7	ND	54.3
092503-01	80.6	80.0	88.2
092503-02	77.9	79.2	81.8
101003-01	83.5	71.6	74.0
101003-02	81.3	69.6	74.3

Surprisingly, these results demonstrate that the stem cell preparations exhibited little or no reduction in viability when stored in syringes on cold packs, thereby demonstrating that stem cell populations may be prepared and shipped, e.g., in a syringe, to a different location for administration to a patient.

EXAMPLE 9

ADIPOSE-DERIVED PLURIPOTENT STEM CELLS FOR TENDON REPAIR

To demonstrate the clinical efficacy of stem cells prepared according to the methods of the invention, as provided in Example 1, a double-blinded, placebo-controlled study was performed using four cell therapy treated and four placebo-treated control horses. Lesions were created with collagenase to mimic natural tendonitis, and autologous cell transplants were conducted 10 days after lesions were induced. Adipose tissue from the tail head area on each animal was processed with the collagenase-based processing protocol. The cell preparations for four of the horses were injected into the lesions, and saline was injected for the controls. Weekly ultrasounds were performed. The horses were sacrificed at six weeks post therapy, and the wound sites examined.

The results of these experiments demonstrate statistically significant improvements in the injuries treated with the stem cell preparations (treated) as compared to the controls. Specifically, inflammatory cell infiltrates were not observed in the treated animals, whereas slight to moderate inflammatory infiltrate was observed in the controls, thus demonstrating the anti-inflammatory effect of the cell preparations. The treated injuries exhibited

significant improvement in architecture with uniform collagen fiber creation and significant improvement in the crimp observed under polarized light indicating normal cross-linking of collagen. In addition, the treated injuries showed statistically significant improvement in the cell treated group demonstrating the overall benefit in improved healing of the tendons. Also, the treated injuries demonstrated improved collagen fiber linearity, more normal tendon shape and sparse tenocyte density, reduced hemorrhage and swelling, more normal new vessel numbers, reduction in lesion size in cells in treated as compared to controls as a percent of starting lesion size (baseline). The lesions in cell treated horses had a lower grade for most of the study as compared to controls, despite starting at an equivalent (or higher) value, and an improved linear fiber pattern was seen.

From the foregoing it will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

CLAIMS

1. A method of preparing a purified cell population comprising stem cells for introducing into a patient, comprising:
 - (a) obtaining adipose tissue from said patient;
 - (b) processing said adipose tissue to separate cells therein from other tissue components; and
 - (c) purifying the separated cells from other tissue components;wherein said method does not include isolating stem cells from other purified cells, thereby preparing a purified cell population comprising adipose tissue-derived stem cells.
2. The method of claim 1, wherein said purified cell population further comprises one or more cells selected from the group consisting of: red blood cells, white blood cells, fibroblasts, fibroblast-like cells, neutrophils, monocyte/macrophages, and basophils.
3. The method of claim 1, wherein said purified cell population further comprises one or more tissue components selected from the group consisting of: extracellular matrix polypeptides or fragments thereof, proteoglycans, cytokines, and growth factors.
4. The method of claim 3, wherein said extracellular matrix polypeptide is selected from the group consisting of: collagen, thrombospondin, fibronectin, vitronectin, cytotactin, laminin and integrins.
5. The method of claim 1, further comprising suspending said purified cells in a physiologically compatible buffer.

6. The method of claim 5, further comprising placing said purified cells in a syringe.

7. The method of claim 1, further comprising freezing said isolated cells in freezing medium.

8. The method of claim 1, wherein said processing comprises one or more procedures selected from the group consisting of:

- (a) mincing the adipose tissue;
- (b) treating the adipose tissue with an enzyme that facilitates the release of cells from other tissue components;
- (c) exposing the adipose tissue to ultrasonic energy; and
- (d) treating the adipose tissue with perfluorocarbons.

9. The method of claim 8, wherein said enzyme treatment is performed at a temperature below 28°C.

10. The method of claim 8, wherein said enzyme treatment is performed at a pH below 7.0.

11. The method of claim 1, wherein said patient is a human.

12. The method of claim 1, wherein said patient is a non-human animal.

13. The method of claim 12, wherein said animal is a horse or camel.

14. The method of claim 12, wherein said animal is a dog or cat.

15. The method of claim 12, wherein said animal is an exotic or zoological animal.

16. The method of claim 15, wherein said animal is a hoofed mammal.

17. The method of claim 15, wherein said animal is a bird.

18. The method of claim 12, wherein said animal is a livestock or farm animal.

19. The method of claim 18, wherein said animal is a cow or goat.

20. The method of claim 12, 13, 16 or 18, wherein said adipose tissue is obtained from the tail head region of the animal.

21. A method of preparing a composition comprising purified adipose tissue-derived stem cells for introducing into an animal, comprising:

(a) obtaining adipose tissue from the tail head region of said animal;

(b) processing said adipose tissue to separate cells therein from other tissue components; and

(c) purifying the separated cells from other tissue components, thereby preparing a composition comprising purified adipose tissue-derived stem cells.

22. The method of claim 21, wherein said composition further comprises one or more cells selected from the group consisting of: red blood

cells, white blood cells, fibroblasts, fibroblast-like cells, neutrophils, monocyte/macrophages, and basophils.

23. The method of claim 21, wherein said composition further comprises one or more tissue components selected from the group consisting of: extracellular matrix polypeptides or fragments thereof, proteoglycans, cytokines, and growth factors.

24. The method of claim 23, wherein said extracellular matrix polypeptide is selected from the group consisting of: collagen, thrombospondin, fibronectin, vitronectin, cytotactin, laminin and integrins.

25. The method of claim 21, further comprising suspending said separated cells in a physiologically compatible buffer.

26. The method of claim 25, further comprising placing said purified cells in a syringe.

27. The method of claim 21, further comprising freezing said isolated cells in freezing medium.

28. The method of claim 21, wherein said patient is selected from the group consisting of: horse and camel.

29. The method of claim 21, wherein said processing comprises one or more procedures selected from the group consisting of:

- (a) mincing the adipose tissue;
- (b) treating the adipose tissue with an enzyme that facilitates the release of cells from other tissue components;
- (c) exposing the adipose tissue to ultrasonic energy; and

(d) treating the adipose tissue with perfluorocarbons.

30. The method of claim 29, wherein said enzyme treatment is performed at a temperature below 28°C.

31. The method of claim 29, wherein said enzyme treatment is performed at a pH below 7.0.

32. A method of providing a composition comprising purified stem cells for introducing into a patient, comprising:

- (a) processing collagen-based tissue obtained from a patient to separate cells therein from other tissue components;
- (b) purifying the separated cells; and
- (c) placing said separated cells in a container, thereby providing a composition comprising purified stem cells.

33. The method of claim 32, further comprising suspending said separated cells in a physiologically compatible solution.

34. The method of claim 32, further comprising shipping said composition to a physician or veterinarian.

35. The method of claim 32, wherein said container is a syringe.

36. The method of claim 32, further comprising freezing or lyophilizing said separated cells.

37. The method of claim 32, wherein said collagen-based tissue is adipose tissue or umbilical cord matrix.

38. The method of claim 32, wherein said method does not include isolating stem cells from other purified cells.

39. The method of claim 32, wherein said purified cell population further comprises one or more cells selected from the group consisting of: red blood cells, white blood cells, fibroblasts, fibroblast-like cells, neutrophils, monocyte/macrophages, and basophils.

40. The method of claim 32, wherein said purified cell population further comprises one or more tissue components selected from the group consisting of: extracellular matrix polypeptides or fragments thereof, proteoglycans, cytokines, and growth factors.

41. The method of claim 40, wherein said extracellular matrix polypeptide is selected from the group consisting of: collagen, thrombospondin, fibronectin, vitronectin, cytotactin, laminin and integrins.

42. The method of claim 32, wherein said processing comprises one or more procedures selected from the group consisting of:

- (a) mincing the tissue;
- (b) treating the tissue with an enzyme that facilitates the release of cells from other tissue components;
- (c) exposing the tissue to ultrasonic energy; and
- (d) treating the tissue with perfluorocarbons.

43. The method of claim 42, wherein said enzyme treatment is performed at a temperature below 28°C.

44. The method of claim 42, wherein said enzyme treatment is performed at a pH below 7.0.

45. The method of claim 32, wherein said patient is a human.
46. The method of claim 32, wherein said patient is a non-human animal.
47. The method of claim 46, wherein said collagen-based tissue is obtained from the tail head region of the animal.
48. A kit useful in the treatment of an injury or disease in an animal, comprising: a container comprising a composition comprising a stem cell population purified from a collagen-based tissue obtained from an animal.
49. The kit of claim 48, wherein said injury is a musculo-skeletal tissue injury.
50. The kit of claim 48, wherein said stem cell population is present in a physiologically compatible solution.
51. The kit of claim 48, wherein said container is a syringe.
52. The kit of claim 48, wherein said container is a cryovial.
53. The kit of claim 48, wherein said composition is frozen.
54. The kit of claim 48, wherein said composition is lyophilized.
55. The kit of claim 48, wherein said collagen-based tissue was obtained from the animal to be treated.

56. The kit of claim 48, wherein said composition further comprises one or more cells selected from the group consisting of: red blood cells, white blood cells, fibroblasts, fibroblast-like cells, neutrophils, monocyte/macrophages, and basophils.

57. The kit of claim 48, wherein said composition further comprises one or more tissue components selected from the group consisting of: extracellular matrix polypeptides or fragments thereof, proteoglycans, cytokines, and growth factors.

58. The kit of claim 57, wherein said extracellular matrix polypeptide is selected from the group consisting of: collagen, thrombospondin, fibronectin, vitronectin, cytotactin, laminin and integrins.

59. The kit of claim 48, wherein said collagen-based tissue is adipose tissue obtained from the tail head region of the animal.

60. The kit of claim 48, wherein said tissue is selected from the group consisting of: tendon, ligament, cartilage, and bone.

61. The kit of claim 48, wherein said tissue is selected from the group consisting of: lung tissue, blood vessels, liver, nerve, and heart.

62. The kit of claim 48, wherein said tissue is hoof laminae.

63. The kit of claim 48, wherein said kit comprises two or more containers, each comprising a composition comprising a stem cell population purified from a collagen-based tissue obtained from an animal.

64. A kit useful in the prevention of an injury in an animal, comprising: a container comprising a composition comprising a stem cell population purified from a collagen-based tissue obtained from an animal.

65. The kit of claim 64, wherein said injury is a musculo-skeletal tissue injury.

66. The kit of claim 64, wherein said stem cell population is present in a physiologically compatible solution.

67. The kit of claim 64, wherein said container is a syringe.

68. The kit of claim 64, wherein said container is a cryovial.

69. The kit of claim 64, wherein said composition is frozen.

70. The kit of claim 64, wherein said composition is lyophilized.

71. The kit of claim 64, wherein said collagen-based tissue was obtained from the animal to be treated.

72. The kit of claim 64, wherein said composition further comprises one or more cells selected from the group consisting of: red blood cells, white blood cells, fibroblasts, fibroblast-like cells, neutrophils, monocyte/macrophages, and basophils.

73. The kit of claim 64, wherein said composition further comprises one or more tissue components selected from the group consisting of: extracellular matrix polypeptides or fragments thereof, proteoglycans, cytokines, and growth factors.

74. The kit of claim 73, wherein said extracellular matrix polypeptide is selected from the group consisting of: collagen, thrombospondin, fibronectin, vitronectin, cytotactin, laminin and integrins.

75. The kit of claim 64, wherein said collagen-based tissue is adipose tissue obtained from the tail head region of the animal.

76. The kit of claim 64, wherein said tissue is selected from the group consisting of: tendon, ligament, cartilage, and bone.

77. The kit of claim 64, wherein said tissue is selected from the group consisting of: lung tissue, blood vessels, liver, nerve, and heart.

78. The kit of claim 64, wherein said tissue is hoof laminae.

79. The kit of claim 64, wherein said kit comprises two or more containers, each comprising a composition comprising a stem cell population purified from a collagen-based tissue obtained from an animal.

80. A method of preparing a purified cell population comprising collagen-based tissue-derived stem cells for delivery to a patient, comprising:

- (a) obtaining collagen-based tissue from said patient;
- (b) processing said collagen-based tissue to separate cells therein from other tissue components; and
- (c) purifying the separated cells,

wherein said processing comprises contacting said collagen-based tissue with a series of screens, thereby preparing a purified cell population comprising collagen-based tissue-derived stem cells.

81. The method of claim 80, wherein said processing further comprises treating the tissue with an enzyme that facilitates the release of cells from other tissue components.

82. The method of claim 80, wherein said purified cell population further comprises one or more cells selected from the group consisting of: red blood cells, white blood cells, fibroblasts, fibroblast-like cells, neutrophils, monocyte/macrophages, and basophils.

83. The method of claim 80, wherein said purified cell population further comprises one or more tissue components selected from the group consisting of: extracellular matrix polypeptides or fragments thereof, proteoglycans, cytokines, and growth factors.

84. The method of claim 83, wherein said extracellular matrix polypeptide is selected from the group consisting of: collagen, thrombospondin, fibronectin, vitronectin, cytostatin, laminin and integrins.

85. The method of claim 80, further comprising suspending said purified cells in a physiologically compatible buffer.

86. The method of claim 85, further comprising placing said purified cells in a syringe.

87. The method of claim 80, further comprising freezing said isolated cells in freezing medium.

88. The method of claim 80, wherein said processing comprises one or more procedures selected from the group consisting of:

(b) exposing the tissue to ultrasonic energy; and

(b) treating the tissue with perfluorocarbons.

89. The method of claim 81, wherein said enzyme treatment is performed at a temperature below 28°C.

90. The method of claim 81, wherein said enzyme treatment is performed at a pH below 7.0.

91. The method of claim 80, wherein said patient is a human.

92. The method of claim 80, wherein said patient is a non-human animal.

93. The method of claim 80, wherein said patient is a horse or camel.

94. The method of claim 80, wherein said patient is a cat or dog.

95. The method of claim 92, wherein said tissue is obtained from the tail head region of the animal.

96. A method of preparing a purified cell population comprising collagen-based tissue-derived stem cells for providing to a patient, comprising:

- (a) obtaining collagen-based tissue from said patient; and
- (b) processing said collagen-based tissue to isolate cells

therein from other tissue components,

wherein said processing comprises contacting the tissue with a surface to which the tissue adheres, thereby preparing a purified cell population comprising collagen-based tissue-derived stem cells.

97. The method of claim 96, wherein said contacting comprises mixing the tissue with particles of the adherent surface.

98. The method of claim 96, wherein said processing further comprises mincing said tissue.

99. The method of claim 98, wherein said processing further comprises treating the tissue with an enzyme that facilitates the release of cells from other tissue components.

100. The method of claim 96, wherein said surface is selected from the group consisting of: Velcro, polystyrene, glass fiber, glass wool, cellulose, and ceramic.

101. The method of claim 96, wherein said purified cell population further comprises one or more cells selected from the group consisting of: red blood cells, white blood cells, fibroblasts, fibroblast-like cells, neutrophils, monocyte/macrophages, and basophils.

102. The method of claim 96, wherein said purified cell population further comprises one or more tissue components selected from the group consisting of: extracellular matrix polypeptides or fragments thereof, proteoglycans, cytokines, and growth factors.

103. The method of claim 102, wherein said extracellular matrix polypeptide is selected from the group consisting of: collagen, thrombospondin, fibronectin vitronectin, cytotactin, laminin and integrins.

104. The method of claim 96, further comprising suspending said purified cells in a physiologically compatible buffer.

105. The method of claim 104, further comprising placing said purified cells in a syringe.

106. The method of claim 96, further comprising freezing said isolated cells in freezing medium.

107. The method of claim 99, wherein said enzyme treatment is performed at a temperature below 28°C.

108. The method of claim 99, wherein said enzyme treatment is performed at a pH below 7.0.

109. The method of claim 96, wherein said patient is a human.

110. The method of claim 96, wherein said patient is a non-human animal.

111. The method of claim 110, wherein said animal is a horse or camel.

112. The method of claim 110, wherein said animal is a dog or cat.

113. The method of claim 110, wherein said adipose tissue is obtained from the tail head region of the animal.

114. A device adapted for preparing a cell population comprising collagen-based tissue-derived stem cells, comprising a series of mesh screens, arrayed such that one or more of the screens may be separated from each other while in contact with a collagen-based tissue sample.

115. The device of claim 114, wherein said screens comprise edges capable of cutting an adipose tissue sample.

116. The device of claim 114, further comprising a cutting implement that may be inserted between adjacent screens.

117. The device of claim 114, further comprising a container comprising said mesh screens.

118. The device of claim 117, wherein said container comprises an opening through which a collagen-based tissue sample may be placed into said container.

119. A method of treating an injury or disease in a patient, comprising providing to said patient an isolated cell population or composition comprising collagen-based tissue-derived stem cells prepared according to a method of any one of claims 1, 21, 32, 80, or 96.

120. The method of claim 119, wherein said injury or disease is a musculo-skeletal injury or disease.

121. The method of claim 119, wherein said isolated cell population or composition is provided directly to a site of injury or disease.

122. The method of claim 119, wherein said isolated cell population or composition is provided by injection.

123. The method of claim 119, wherein said isolated cell population or composition is provided to the bloodstream of said patient.

124. The method of claim 119, wherein said isolated cell population or composition is provided intravenously or inter-arterially.

125. The method of claim 119, wherein said tissue is selected from the group consisting of: tendon, ligament, cartilage, and bone.

126. The method of claim 119, wherein said tissue is hoof laminae.

127. The method of claim 119, wherein said tissue is selected from the group consisting of: lung, blood vessels, liver, nerve, and heart.

128. The method of claim 119, wherein said injury is selected from the group consisting of: sprain, strain, dislocation, bruising, tear, and fracture.

129. The method of claim 119, wherein said injury or disease is an ischemic injury or disease.

130. The method of claim 119, wherein said injury or disease is a septic injury or disease.

131. The method of claim 119, wherein said patient is a human.

132. The method of claim 119, wherein said patient is a non-human animal.

133. The method of claim 119, wherein said animal is a horse or camel.

134. The method of claim 119, wherein said animal is a dog or cat.

135. A method of preventing an injury in a patient, comprising providing to said patient an isolated cell population or composition comprising collagen-based tissue-derived stem cells prepared according to a method of any one of claims 1, 21, 32, 80, or 96.

136. The method of claim 135, wherein said injury is a musculo-skeletal injury.

137. The method of claim 135, wherein said isolated cell population or composition is provided directly to a site of potential injury.

138. The method of claim 135, wherein said isolated cell population or composition is provided by injection.

139. The method of claim 135, wherein said isolated cell population or composition is provided to the bloodstream of said patient.

140. The method of claim 135, wherein said isolated cell population or composition is provided intravenously or inter-arterially.

141. The method of claim 135, wherein said tissue is selected from the group consisting of: tendon, ligament, cartilage, and bone.

142. The method of claim 135, wherein said tissue is hoof laminae.

143. The method of claim 135, wherein said tissue is selected from the group consisting of: lung, blood vessels, liver, nerve, and heart.

144. The method of claim 135, wherein said injury is selected from the group consisting of: sprain, strain, dislocation, bruising, tear, and fracture.

145. The method of claim 135, wherein said injury or disease is an ischemic injury or disease.

146. The method of claim 135, wherein said injury or disease is a septic injury or disease.

147. The method of claim 135, wherein said patient is a human.

148. The method of claim 135, wherein said patient is a non-human animal.

149. The method of claim 135, wherein said animal is a horse or camel.

150. The method of claim 135, wherein said animal is a dog or cat.